





Managing Sick Workers Best Practices Act Immediately Protect the person handling sick workers Have an action plan ready Isolation for quarantining Medical treatment Access to medical support

13

Managing Sick Workers Best Practices ✓ Prevent spread → Contact Tracing (2-days, 6-feet and 10sec.) → Notify and quarantine known contacts → Educate workers living off property → Effecting training to understand risk controls → Clean, disinfect and PPE use

14

Managing Sick Workers Return to Work — Persons Under COVID-19 Isolation 1) Symptom-based strategy (Positive with symptoms) >3 days (72 hours) — No fever, no cold medications, no respiratory symptoms; and, >10 days have passed since symptoms first appeared. 3) Time-based strategy (Positive without symptoms) >10 days since positive test, free from any symptoms. 2) Test-based strategy > Resolution of fever without the use of fever-reducing medications and > Improvement in respiratory symptoms (e.g., cough, shortness of breath), and > Two consecutive test ≥ 24-hour period







Best Practices Provide training and guidance of risk controls Common area cleaning and disinfecting Employer and employee disinfecting policies Use proper cleaning chemicals Dormitory style housing need to be quartered into smaller groups Keep family members together

19



20









MEMIC ©
























MEMIC © 11







Pandemic Safety for the Migrant Worker 1) Risk Assessment and Control Plan 2) Arrival and Screening 3) Managing a Sick Worker 4) Shared Housing 5) Transportation 6) Worksite & Work Areas